



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

curator; they both agreed that the bird should be regarded as an Essex County record. According to Mr. Townsend's 'Birds of Essex County,' and Howe and Allen's 'Birds of Massachusetts,' this is the first authentic record of the occurrence of the Clapper Rail in Essex County, though both cite: "Mr. J. F. Le Baron informed me that he shot a specimen some years ago at Ipswich." C. J. Maynard, the Naturalist's Guide, p. 145, 1870.

The mounted bird will be presented to the Peabody Museum, Salem.—
WILLIAM P. WHARTON, *Groton, Mass.*

Late Flight of Woodcock on Long Island, N. Y.—On December 5, 1908, while hunting near Flanders, Suffolk Co., I flushed a Woodcock (*Philohela minor*). On December 8, after a storm and heavy rain, eleven were shot in a small swamp at Lawrence, Nassau Co., close to the New York City line. The same day one was seen in a little strip of woods about one mile from this swamp.

In 30 years on Long Island, I have never seen other than stragglers after about November 20th and do not recall ever before seeing one in December.—HAROLD HERRICK, *New York City.*

Capture of the Ruff at Seabrook, N. H.—Mr. John Hardy of Boston has kindly presented me with an adult female Ruff (*Pavoncella pugnax*) shot at Seabrook, N. H., Sept. 23, 1907, by Charles Fowler, who said it was with a flock of Black-bellied Plover.—JOHN E. THAYER, *Lancaster, Mass.*

Eskimo Curlew taken at Newburyport, Mass.—I purchased of Mr. John Hardy of the Boston Market, a male Eskimo Curlew (*Numenius borealis*) taken at Newburyport, Mass., by A. B. Thomas, August 27, 1908. He shot two, but the other bird had its head so badly shot that it could not be made into a skin.—JOHN E. THAYER, *Lancaster, Mass.*

The American Golden Plover (*Charadrius dominicus*) in Ohio in Autumn.—On October 23, 1908, I met a flock of 6 American Golden Plovers at the Grand Reservoir, Ohio, and on October 27 a pair of them was shot, together with a Baird's Sandpiper and a pair of Wilson's Snipe at the Loramie Reservoir in Shelby Co., O. Fall records in Ohio for the American Golden Plover are very rare, in fact any record of the occurrence of this species in this State is interesting and noteworthy. The female of this pair of Golden Plovers has the tip of the upper mandible curved over the lower one to the extent of about 4 mm., with the tip curving decidedly downward almost at right angles, thus forming a veritable crossbill. The cause of this formation cannot be seen, only the upper mandible seems to be rather thin and weak, when compared with that of the other specimen, nor was the bird as fat as the male.—W. F. HENNINGER, *New Bremen, Ohio.*